

Occurrence of ESBL-producing *Escherichia coli* ST131 including the H30-Rx and C1-M27 subclones amongst urban seagulls from UK

Flavia Zendri¹*, Iuliana E. Maciuca¹*, Simon Moon², Philip H. Jones³, Andy Wattret¹, Richard Jenkins⁴, Andy Baxter⁵, Dorina Timofte^{*1}

¹ Institute of Veterinary Science, University of Liverpool, UK; ² Somerset West and Taunton Council, Taunton, UK; ³ Animal and Plant Health Agency (APHA), Shrewsbury, UK; ⁴ School of Allied Health Sciences, De Montfort University, Leicester, UK; ⁵ Birdstrike Management Ltd., York, UK;

© These authors contributed equally

*Corresponding author:

Dorina Timofte

Institute of Veterinary Science

Department of Veterinary Pathology and Public Health

Faculty of Health & Life Sciences

University of Liverpool, Leahurst Campus

Chester High Road

CH64 7TE, Neston

England

E-mail: d.timofte@liv.ac.uk

Phone no: (0151) 794 6193

Running title: ESBL-producing *E. coli* among urban UK seagulls

Abstract

Antimicrobial resistance is a public health concern. Understanding any role that urban seagulls may have as a reservoir of resistant bacteria could be important for reducing transmission.

This study investigated faecal *Escherichia coli* isolates from seagulls (herring gulls and lesser black-backed gulls) to determine the prevalence of extended-spectrum cephalosporin resistant (ESC-R) and fluoroquinolone-resistant (FQ-R) *E. coli* among gull species from two cities (Taunton and Birmingham) in the United Kingdom. We characterised the genetic background and carriage of plasmid-mediated resistance genes in extended spectrum β -lactamase (ESBL)-producing *E. coli* obtained from these birds.

Sixty ESC-R *E. coli* isolates were obtained from 39 seagulls (39/78, 50%), of which 28 (28/60, 46.7%) were positive for plasmid-mediated CTX-M and/or AmpC β -lactamase resistance genes. Among these, *bla*_{CTX-M-15}, *bla*_{CTX-M-14} and *bla*_{CMY-2} predominated. Three isolates belonging to the B2-ST131 clone were detected, of which two harboured *bla*_{CTX-M-15} (typed to C2/H30Rx) and one harboured *bla*_{CTX-M-27} and was typed to C1/H30-R (recently described as the C1-M27 sublineage). PMQR gene carriage prevalence (11.7%) consisted of *aac(6')-Ib-cr* and *qnrB* genes. No carbapenem or colistin resistance genes were detected.

Urban seagulls in the UK are colonised and can spread major antimicrobial resistant *E. coli* isolates harbouring ESBL and PMQR determinants, including clinically important strains such as the pandemic clone B2-ST131 and the C1-M27 subclade. This is the first report of ST131-C1-M27 subclade in wildlife in the UK and in seagulls worldwide.

Keywords: *Escherichia coli*, B2-ST131, *Larus* spp., ESBLs, AmpC, PMQR, C1-M27

1. Introduction

In the past two decades, extra-intestinal pathogenic *Escherichia coli* (ExPEC) has been the leading cause of difficult-to-treat human urinary tract infections (UTI), soft tissue and bloodstream infections, in both healthcare and community-onset settings.¹ Selection pressure from antibiotic use has shaped the rapid evolution and spread of antimicrobial resistance among *E. coli* strains no longer responding to conventional treatment and also to critically important antimicrobials such as 3rd generation cephalosporins.² Antimicrobial resistance appears concentrated within specific bacterial genotypes, among which *E. coli* sequence type (ST) 131, with specific reference to the O25b:H4 serotype, has received particular attention owing to its rapid emergence and pandemic global spread.^{3,4} Although the reasons behind the successful dissemination of *E. coli* B2-ST131-O25b:H4 have not been entirely elucidated, a large number of virulence-related genes combined with transferable antimicrobial resistance determinants carried on plasmids have been identified as possible factors.⁵⁻¹⁰ In addition, *E. coli* ST131 isolates have been contributing to the recent CTX-M spread, leading to a global rise in *Enterobacteriaceae* harbouring CTX-M type extended-spectrum β -lactamases (ESBLs).¹¹ There is now increasing evidence that the environment is a vast reservoir of resistant bacteria and their associated genes and that complex interactions exist between the environmental resistome and that of clinically significant pathogens.^{12,13} In this context, current evidence suggests that wild birds, including seagulls, are contributing to the environmental dissemination of antibiotic resistance,¹⁴ which is of great concern when considering highly pathogenic clones such as *E. coli* ST131.

Population genetic studies have given a new insight into the clonal structure of *E. coli* ST131 by revealing epidemiologically distinct sublineages nested within ST131 clade C/H30.¹⁵ Clade C is known for expressing high-level ciprofloxacin-resistance mutations resulting in distinctive *gyrA* and *parC* variants^{16,17} and, nowadays, it accounts for the vast majority of *E. coli* ST131

worldwide.¹⁸ Whole-genome sequencing (WGS) enabled the characterisation of the two main subclades within C/H30, namely C1/H30R (non-RX) and C2/H30Rx.^{19,20} The latter is a clonal subset often associated with *bla*_{CTX-M-15}, whose clonal expansion was largely responsible for the *E. coli* ST131 pandemic and, along with it, to the substantial global rise in CTX-M-15 (currently the most widespread CTX-M-type ESBL enzyme in humans and animals worldwide).^{17,21–23} Conversely, isolates belonging to the C1/H30R subclade are largely ESBL negative, although certain isolates harbour *bla*_{CTX-M-14} or *bla*_{CTX-M-27}, the former being the second most common ESBL type occurring in humans globally whereas the latter is an ESBL type more often recovered among animal and environmental ST131 strains.²⁴

Notwithstanding *E. coli* ST131 being a major cause of disease in people, a few reports have described the detection of pandemic clone B2-ST131 carrying diverse CTX-M-type enzymes in animals, foodstuffs, and the environment.^{3,25–30}

Although there are high-level similarities between certain human and animal ST131 isolates with regard to resistance characteristics, virulence factors and genetic background, there is only limited evidence for direct inter-species transfer of ST131.³¹ Given the importance of ST131 as a global public health issue, further investigation of animal-associated ST131 is warranted in order to clarify the ecology and capacity for cross-species transmission of these strains.

Wild birds have been proposed as sentinels and potential vectors of antimicrobial resistance since many species, such as geese, mallards, cormorants, pigeons, corvids, gulls and birds of prey, have been found carriers of diverse multi-drug resistant (MDR) bacteria including *E. coli* strains resistant to extended-spectrum cephalosporins (ESCs) and fluoroquinolone (FQs).^{32–41} In this regard, members of the *Laridae* family of seabirds have become particularly studied as sentinels for monitoring the spread of antimicrobial resistance since their ecology is well understood and they possess several characteristics which make them suited to dissemination

studies.³⁸ Furthermore, gull species are regarded as important bio-indicators of environmental contamination by antibiotic resistance, with particular regard to surface waters and coastal areas.⁴²⁻⁴⁴ In addition, gull species can migrate over long distances and they have proven capable of introducing MDR *E. coli* in remote areas dominated by ecosystems with little or no antibiotic resistance, as illustrated by the recovery of CTX-M-15 producing *E. coli* strain belonging to the pandemic clone B2-ST131 in the Russian Commander Islands.⁴⁵

The aim of this study was dual; firstly, to determine the faecal prevalence of resistance to ESCs amongst seagull species from two cities in the United Kingdom (one coastal and one inland) and secondly, to characterise the genetic background as well as plasmid-mediated resistance genes, virulence genes and plasmid incompatibility groups in ESBL-producing *E. coli* obtained from these birds.

2. Materials and Methods

2.1 Collection of faecal samples

Herring gull (*Larus argentatus*) and lesser black-backed gull (*Larus fuscus*) faecal samples were collected from two locations (L1: Taunton and L2: Birmingham) in England, United Kingdom, as part of Local Authority non-lethal pest-control programmes designed to minimise public nuisance caused by the gulls. The method used involved replacing gull eggs in the nests with decoy eggs that encouraged gulls to remain on the nests and minimised the public nuisance caused by gulls scavenging for food within the cities. In response to being disturbed, gulls frequently produced a faecal sample that was collected into sterile universal tubes. Sampling took place in May 2011 over a two days' period. For L1, 50 faecal samples were collected on the first day in the town centre from $n=39$ herring gulls and $n=11$ lesser black-backed gulls. For L2, 28 faecal samples were collected on the second day in a single district of the city from

$n=7$ herring gulls and $n=21$ lesser black-backed gulls. Upon collection, samples were placed in sterile Universal tubes and transported rapidly to the laboratory for immediate processing; when this was not possible, samples were refrigerated for next-day processing.

2.2 Bacterial isolates

Faecal samples (1-2 g) were placed in 10 ml of Buffered Peptone Water (BPW, Thermo Scientific, Basingstoke, UK) and incubated overnight at 37°C.

To screen for ESC-resistant *E. coli*, a small BPW inoculum (2-3 µl) was streaked onto eosin methylene blue agar (EMBA, Thermo Scientific, Basingstoke, UK) containing 1 µg/ml of cefotaxime (all antibiotics from Sigma-Aldrich Ltd., UK). If colonies were phenotypically different, each colony morphotype was selected.

2.3 Antimicrobial susceptibility testing

Any cefotaxime-resistant isolates obtained on the EMBA selective media, producing a characteristic *E. coli* green metallic sheen (and subsequently confirmed as *E. coli* by *uidA* and *uspA* polymerase chain reaction [PCR] assay) and for which plasmid-mediated ESC and/or FQ resistance was identified, were subjected to antimicrobial susceptibility testing. The susceptibility testing was performed by disk diffusion on Mueller-Hinton agar (MHA) according to the Clinical and Laboratory Standards Institute methodology (CLSI) 2018.⁴⁶ *E. coli* ATCC25922 was used as control for disk diffusion susceptibility testing for every new batch of isolates using a suspension of 0.5 McFarland to obtain a semiconfluent lawn on MHA. The antimicrobial panel comprised ampicillin (10 µg), amoxicillin/clavulanic acid (30 µg), cefoxitin (30 µg), cefpodoxime (10 µg), cefepime (30 µg), piperacillin/tazobactam (110 µg), aztreonam (30 µg), imipenem (10 µg), ertapenem (10 µg), nalidixic acid (30 µg), gentamicin (10 µg), amikacin (30 µg), streptomycin (10 µg), chloramphenicol (30 µg), doxycycline (30 µg) and trimethoprim/sulfamethoxazole (25 µg) (all disks and media from Oxoid, UK).

Interpretation of results was performed according to the Clinical and Laboratory Standards Institute criteria (CLSI) 2018.⁴⁶

2.4 Characterisation of resistance genes

In order to identify the resistance genes carried by the ESC-resistant *E. coli* isolates, cell lysates were prepared by suspending the bacterial cells in 500 µl sterile distilled water followed by incubation at 95°C for 10 minutes and centrifugation at 10,000 g for 5 minutes. All ESC-resistant isolates were subjected to PCR screening for the presence of *bla*_{SHV}, *bla*_{TEM}, *bla*_{OXA1-like}, and *bla*_{CTX-M} genes, with a further multiplex PCR being used for the detection of family-specific plasmid-mediated AmpC β-lactamase genes.^{47,48} Furthermore, plasmid-mediated quinolone resistance (PMQR) genes *qnrA*, *qnrB*, and *qnrS*, *qepA*, and the *aac(6')-ib-cr* gene, which confers reduced susceptibility to aminoglycosides and fluoroquinolones, were identified.^{49,50} All isolates positive for *aac(6')-ib* gene were sequenced with specific primers to confirm the presence of *aac(6')-ib-cr* variant.⁴⁹ Additionally, genetic determinants encoding for resistance to carbapenems and colistin, were investigated according to the methods described by Dallene *et al.*⁴⁷, Poirel *et al.*⁵¹, Liu *et al.*⁵² and Rebelo *et al.*⁵³ To confirm gene identity, DNA sequencing of both strands of PCR reaction products was done for representative isolates from groups displaying unique gene combinations (e.g., *bla*_{CTX-M}, *bla*_{TEM}, *aac(6')-ib-cr*) using the same sets of primers as in the original reactions (Eurofins MWG Operon). The resulting DNA sequences were compared using BLASTn against sequences in GenBank. All ESC-R *E. coli* isolates were selected for further molecular typing.

2.5 Characterisation of virulence genes

ESC-R *E. coli* seagull isolates were screened by multiplex PCR for virulence determinants associated to diarrheagenic Shiga-toxin producing (STEC) strains, including Shiga-like toxins 1 and 2 (encoded by *stx*₁ and *stx*₂) and for accessory STEC virulence factors intimin (*eaeA*) and

enterohaemorrhagic (EHEC) plasmid-mediated enterohaemolysin (*hlyA*) according to Paton and Paton⁵⁴ and Vidal *et al.*⁵⁵ EHEC belonging to serotypes displaying greater virulence, i.e. O157 and O111, were investigated by multiplex PCR assay as previously described.⁵⁴ Furthermore, PCR targeting of virulence genes linked to uropathogenicity, including the cytotoxic necrotizing factor type 1 and 2 (*cnf1* and 2) and various classes of adhesins, such as the pyelonephritis-associated pili (*pap*), S-fimbrial adhesin (*sfa*) and afimbrial adhesin (*afa*), was performed as described by Blanco *et al.*⁵⁶

2.6 Molecular characterisation of the ESC-R *E. coli* isolates

E. coli isolates were assigned to phylogenetic groups (A, B1, B2, C, D, E, F and clade I) by multiplex PCR using two methods, the triplex and the revisited quadruplex methods proposed by Clermont *et al.*^{57,58} The isolates identified as phylogenetic group B2 were screened for the O25 group by a method based on an allele-specific PCR.⁵⁹ Positive isolates were further tested to identify members of the international clone B2-ST131 using primers for the genes *pabB* and *trpA*, as described by Clermont *et al.*⁶⁰ Following screening for the B2-ST131 clone, positive isolates carrying *bla*_{CTX-M-27}, *bla*_{CTX-M-14} and *bla*_{CTX-M-15} genes underwent further analysis consisting of multiplex conventional PCR in order to detect all ST131 clades (A, B and C) as well as C subclades (C1-M27, C1-nM27 [C1-nonM27] and C2), as developed by Matsumura *et al.*⁶¹

2.7 Resistance transfer and PCR-based replicon typing

Transfer of resistance genes by conjugation was performed by broth mating to determine whether the ESBL resistance determinants were transferable. The conjugation assays were attempted for isolates carrying *bla*_{CTX-M} genes using a streptomycin-resistant *E. coli* HB101 strain as a recipient as previously described by Oliver *et al.*⁶² Transconjugants were selected on nutrient agar (Oxoid) supplemented with streptomycin (50 µg/ml) and cefotaxime (1 µg/ml). The resistance phenotype of transconjugants was determined by disk diffusion using the same

antimicrobial panel as for parental strains. Plasmid replicons were determined on all *bla*_{CTX-M} positive isolates and transconjugants, using a PCR-based replicon-typing scheme as previously described.⁶³

3. Results

3.1 Bacterial isolates and antimicrobial susceptibility testing

Screening of the 78 seagull faecal samples collected from both locations identified 60 ESC-R *E. coli* isolates obtained from 39 faecal samples, resulting in 50% ESC-R *E. coli* carriage in the seagull population investigated; fifty-five ESC-R *E. coli* isolates were obtained from L1 (from 34 seagulls) and five isolates from L2 (from five seagulls). Antimicrobial susceptibility testing showed high levels of resistance to ampicillin (100%), cefpodoxime (100%), aztreonam (91.6%), amoxicillin/clavulanic acid (83.8%), streptomycin (70.3%), cefoxitin (62.2%), tetracycline (56.8%), and trimethoprim/sulfamethoxazole (51.4%). In addition, 10.8% and 5.4% of the *E. coli* isolates were resistant to ertapenem and imipenem, respectively.

3.2 Characterisation of resistance and virulence genes

Screening for ESBL and AmpC genetic determinants identified 46.7% (28/60) of isolates positive for plasmid-mediated CTX-M and/or AmpC β -lactamase resistance genes. The *bla*_{CTX-M} genes were identified in 21/60 (35%) of the seagull ESC-R *E. coli* isolates. Of these 21, 12 belonged to CTX-M group 1 (57.1%) and sequencing identified *bla*_{CTX-M-15} to be the most prevalent resistance gene carried by eleven isolates (91.7%) while only one isolate harboured *bla*_{CTX-M-1}. Also nine of the 21 *E. coli* *bla*_{CTX-M} positive isolates (42.9%) carried CTX-M group 9, with six of the nine harbouring *bla*_{CTX-M-14} (66.7%) and *bla*_{CTX-M-27} being present in the remaining three isolates (33.3%). Resistance genes encoding for CTX-M group 2, 8 and 25 were not

identified. It is important to note that none of the investigated seagull isolates harboured the *bla*_{SHV} gene.

A variety of gene combinations was found in the CTX-M-producing *E. coli* isolates with *bla*_{CTX-M-15}, *bla*_{CTX-M-14} and *bla*_{CTX-M-27} carried alone or in combination with *bla*_{TEM}, *bla*_{OXA1-like} and *aac(6')-ib-cr* gene (Table 1). When screened for AmpC type β -lactamase, 13.3% (8/60) of seagull ESC-R *E. coli* isolates carried *bla*_{AmpC} resistance genes, borne singularly or in various combinations with *bla*_{CTX-M-15}, *bla*_{TEM}, *bla*_{OXA1-like}, *aac(6')-ib-cr* and *qnrB*. Sequencing identified all *bla*_{AmpC} genes as *bla*_{CMY-2}. The overall PMQR genes prevalence among seagull ESC-R *E. coli* isolates was 11.7% (7/60) and these consisted of five *aac(6')-ib-cr* and three *qnrB* positive strains, one of which co-harboured both determinants. None of the investigated isolates carried carbapenem resistance genes nor *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* determinants conferring resistance to colistin.

A PCR assay targeting STEC-associated *stx1* and *stx2* genes and *E. coli* serotypes O111 and O157 was negative for all *E. coli* seagull isolates. One isolate carried the gene encoding for intimin (*eaeA*) whilst none was positive for plasmid-encoded enterohaemolysin (*hlyA*). The vast majority of uropathogenic genes were carried individually in *E. coli* seagull isolates, with just two virulence gene combinations occurring in four isolates (Table 1). Of the 23 *E. coli* isolates carrying urovirulence determinants (mainly *pap1/2*⁺), 78.2% (18/23) also harboured AMR genes whilst 21.8% (5/23) were negative for any of the AMR genes tested.

3.3 Molecular typing of ESC-R *E. coli*

All *E. coli* seagull isolates initially assigned to phylogenetic groups using the triplex PCR method, were subsequently re-tested using the extended quadruplex approach. Whilst there was consistency for allocation of the isolates within PGs B1, B2 and D between the two methods, the quadruplex method allowed to redistribute a substantial fraction of the isolates formerly

ascribed to phylogroup A into the later recognized phylogroups C, E, F and clade I as follows: A ($n=4$), B1 ($n=18$), B2 ($n=12$), C ($n=2$), D ($n=12$), E ($n=1$), F ($n=4$), clade I ($n=2$) and unassignable ($n=5$). Overall, phylogroup B1 was the most represented among the seagull isolates (30%); nonetheless, the proportion of isolates belonging to phylogroup B2, which are more commonly associated with extra-intestinal clinical infection,^{64,65} appears high (20%). Furthermore, our study indicates that approximately 36% and 21% of all ESBL isolates ($n=28$) belonged to D and B2 phylogroups, respectively, while both phylogroups B1 and F accounted for 14% of all ESBL seagull isolates. Among the seagull *E. coli* isolates belonging to phylogroup B2, three (25%) have been typed to the international clone B2-ST131-O25b based on the detection of *pabB* and *trpA* genes. This was subsequently confirmed using the multiplex conventional PCR targeting all ST131 clades and the C subclades which typed two of the isolates carrying *bla*_{CTX-M-15} to the C2/H30Rx subclade whilst one isolate carrying *bla*_{CTX-M-27} belonged to the C1-M27 subclade of ST131.

3.5 Resistance transfer and PCR-based replicon typing

Conjugation experiments demonstrated transferability of *bla*_{CTX-M-15} as well as of *bla*_{CTX-M-1} and *bla*_{CTX-M-27} to the *E. coli* recipient in seagull isolates. In addition, PCR-based replicon typing (PBRT) showed that IncFIA and Y were associated with transfer of ESBL genes. IncFIA was the most common replicon found in the seagull *E. coli* transconjugants, being recovered from four out of six transconjugants in which the CTX-M genes were successfully transferred. In the case of one *bla*_{CTX-M-15} transconjugant, the recovered plasmids were not typeable using PBRT whilst for *bla*_{CTX-M-1} the transconjugant obtained lacked IncFIA identified in the original isolate. For *bla*_{CTX-M-27}, both IncFIA and Y replicon types were recovered from the single successful transconjugant. None of the *bla*_{CTX-M-14} genes were transferred, indicating that they may be located on a nonconjugative plasmid or on the chromosome.

4. Discussion

The results described in this study provide further evidence that wild seagulls in Europe may be a reservoir and contribute to dissemination of ESC- and F-resistant *E. coli*. This is in agreement with previous studies indicating the ability of gulls to carry and disseminate *E. coli* strains harbouring several TEM and CTX-M types extended-spectrum β -lactamase^{41,66-72} alone or combined with AmpC β -lactamase^{73,74} and, more recently, with VIM type carbapenemase.^{75,76}

Here, we report a high prevalence (50%) of ESC-R *E. coli* carriage in the UK urban seagull population investigated, amongst which *bla*_{CTX-M-15} was the predominant ESBL genotype. This is much higher than the ESC-R *E. coli* carriage identified in a recent study in silver gulls from Australia (21.7%)⁷⁷ or amongst North American gulls and pelicans (14%)⁷³ and European gulls (28.7%).⁷² In the European study, ESBL-producing *E. coli* from English gulls were detected at a prevalence of 27.1%, with Spain only displaying higher prevalence than the UK (74.8%).⁷²

Consequently, seagulls can act as sentinels and bio-indicators of antimicrobial-resistance and should be regarded as vehicles and potential long-term reservoirs of multidrug-resistant bacteria. Current global antimicrobial drug resistance data indicate that environmental reservoirs are likely to contribute to the selection and spread of resistance determinants and antimicrobial-resistant bacteria.⁷⁸ Despite not being directly exposed to clinically-employed antimicrobial agents, wild animals can be an underestimated vector of antimicrobial resistance as they may acquire it through contact with human-influenced habitats of any sort, such as manure-fertilized meadows, livestock farms, landfills, hospital sewage systems, wastewater treatment facilities, recreational waters and others.⁷⁹ Their feeding behavior and association with water environments are epidemiologically relevant for the gulls' capacity to spread antibiotic-resistance and both coastal and urban environments provide potentially diverse

sources of food. Furthermore, levels of resistance seem to correlate with the extent of association to human activities.⁸⁰ A few studies have shown gulls' uptake of MDR *E. coli* from human waste and sewage facilities (including ST131 strains and, notably, one *mcr-1* positive isolate)⁸¹⁻⁸³ and their ability to acquire antibiotic resistance from these sources and release it into beach waters or other aquatic environment.^{71,74,84,85} **This is particularly relevant for transmission since gulls are migratory birds and can convey antimicrobial resistant bacteria and associated genes into distant environments and populations, including humans, interconnecting environmental niches of different antimicrobial resistance levels.** Moreover, clusters of genetically comparable ESBL producing *E. coli* have been simultaneously isolated from humans and gulls in France⁶⁷, Sweden⁸⁶ and Chile⁸⁷, indicating that transfer between human populations and the environment, including wild birds, is likely to occur. One Alaskan study described the increased detection of MDR *E. coli* from gulls specifically in urban areas rather than rural ones.⁸⁸

The pandemic virulent *E. coli* B2-ST131 clone was rare among the ESC-R *E. coli* isolates analyzed (5%), with 2/3 isolates harbouring *bla*_{CTX-M-15} whilst 1/3 harboured *bla*_{CTX-M-27}. B2-ST131 has been widely shown to be an important driver of the increase in antimicrobial resistance and to have contributed to the dissemination of the CTX-M β -lactamase worldwide.¹⁵ Recent epidemiologic and genomic studies have shown that particular *E. coli* ST131 subclades such as the C2/H30Rx have been largely responsible for the dissemination of CTX-M-15 β -lactamase worldwide¹⁹ whilst the C1/H30R (recently described as the C1-M27) was involved in the epidemic spread of ST131-CTX-M-27 in Asia.²⁴ This distinct *E. coli* ST131 subclade, carrying *bla*_{CTX-M-27} and defined by unique genomic characteristics, has recently emerged within C1/H30R and has rapidly established as the predominant ST131 type and most important vehicle of ESBL-producing ExPEC in Japan.⁸⁹ Following its recognition in other Asian countries,⁹⁰ this novel ST131 subclade, now commonly referred to as C1-M27, has been

increasingly reported also in Europe, with some countries, such as France and Switzerland, having disclosed alarmingly high prevalence rates.⁹¹⁻⁹⁴ Little data is available regarding the presence of *E. coli* ST131 C1-M27 in the veterinary field. Sporadic reports of *E. coli* ST131 carrying *bla*_{CTX-M-27} in animals include clinical specimens from dogs and cats, mostly suffering from UTIs, in Japan and Europe⁹⁵⁻⁹⁹ and faecal samples from wild birds, namely two great cormorants, identified in the Czech Republic³⁹ and Switzerland.¹⁰⁰ CTX-M-27 producing ST131 isolates have also been detected in diverse aquatic environments in Asia and Europe, including hospital wastewater treatment plants,¹⁰¹ rivers and lakes^{102,103} and fish.²⁷ Furthermore, a recent study from the Czech Republic has demonstrated that ST131 harbouring *bla*_{CTX-M-27} from different sources including wild water birds, wastewater and hospital isolates, shared extensive genetic commonalities between them and with isolates previously reported from Japanese hospitals.²⁶ To date, coastal and river waters in the UK have been found polluted with B2-ST131 *E. coli* producing CTX-M-15 and CTX-M-14 enzymes but not CTX-M-27.¹⁰⁴⁻¹⁰⁶ To the best of our knowledge, this is the first report of an *E. coli* ST131 strain belonging to C1-M27 in seagulls and in UK wildlife.

In our study, the CTX-M-15 type enzyme was also prevalent among non-ST131 ESBL-producing *E. coli*, followed by CTX-M-14 type. This is in agreement with previous findings of non-ST131 *E. coli* CTX-M-15 and CTX-M-14 producing ESBL types being frequently isolated from wild gulls (and other wild bird species).⁴¹ Conversely, we report a lower prevalence of *bla*_{CTX-M-1} among non-ST131 ESBL-producing *E. coli* in our UK seagull population than was previously described by other studies.⁷²

As prevalence rates of *E. coli* ST131 in people appear consistently higher than those found in animals in diverse geographic areas, it is generally presumed that humans are more important hosts of ST131 clinical strains compared with animals^{3,107} and that spill-over has later occurred into the veterinary field. Consistent distribution of ESBL genotypes, including ST131, has

generally been described in humans whereas that observed in animals appears more diversified and incongruent (also geographically), with partial overlap of human and animal genotypes. In contrast to domesticated animals, the presence and epidemiology of ESC-R and FQ-R *E. coli*, including medically important and globally widespread clonal lineages like B2-O25b-ST131, have been infrequently addressed in wildlife and the extent and pathways through which these pathogenic bacteria are spread between different ecological niches remain to be elucidated.

In the present study, seagulls colonised with ESBL *E. coli*, including B2-ST131 isolates, were associated with urban areas in a coastal city and we presume they might have acquired them from those environments. The countryside surrounding Taunton stretches up to the coast and offers numerous water bodies and reservoirs serving as overnight roosting sites as well as various agricultural food sources whilst Birmingham is a much larger and populated inland city, placed at greater distance from water bodies but with abundance of landfills and establishments to provide urban gulls with food. The reasons behind the lack of isolation of ESBL-producing *E. coli* among seagulls sampled in the inland city could be partially explained by the smaller sample size or it may be that coastal areas in the UK provide greater opportunities for seagulls to become colonised than those offered by urban environments without close proximity to waterfronts. One limitation of this study comprise the small scale and lack of a far-reaching geographical systemic approach and the absence of parallel investigations into human or environmental ESBL *E. coli*, including ST131-C1-M27 strains, in the same areas as well as the possible relatedness of our C1-M27 isolate with Asian reference strains.

Conclusions

Wild seagulls in the UK are colonised, and therefore can spread, major antimicrobial resistant *E. coli* isolates including clinically important strains such as the pandemic B2-ST131 clone,

and the emerging C2/H30Rx and the C1-M27 subclades. To the best of our knowledge, this is the first report of the ST131-C1-M27 in wildlife in the UK and in seagulls worldwide. Findings from this study support the concept that carriage of these multidrug-resistant strains is widespread among wild gulls, at least in some countries. Further studies are needed to explore the implications of such strains carriage by wild seabirds and to unravel the potential for ongoing interspecies transmission between humans and urban opportunistic birds. As available evidence is pointing towards humans as the original source for ESBL, AmpC and PMQR enrichment of *E. coli* strains in wild species, priority must be given to break the transmission chain between humans and the environment, including wildlife, to prevent further dissemination of MDR bacteria at global level through these birds' migrations.

Disclosure Statement

The authors have no competing financial interests/conflicts to declare.

Funding

This work was funded by the Institute of Veterinary Science, University of Liverpool.

References

1. Dale, A.P., and N. Woodford. 2015. Extra-intestinal pathogenic *Escherichia coli* (ExPEC): disease, carriage and clones. *J. Infect.* 71:615–626.
2. Carvalho, I., N. Silva, J. Carrola, V. Silva, C. Currie, G. Igrejas, and P. Poeta. 2019. Antibiotic Resistance: Immunity-Acquired Resistance: Evolution of Antimicrobial Resistance Among Extended-Spectrum β -Lactamases and Carbapenemases in *Klebsiella pneumoniae* and *Escherichia coli*. In *Antibiotic Drug Resistance*. John Wiley & Sons, Ltd, pp. 239–259.

3. Nicolas-Chanoine, M.-H., X. Bertrand, and J.-Y. Madec. 2014. *Escherichia coli* ST131, an Intriguing Clonal Group. *Clin. Microbiol. Rev.* 27:543–574.
4. Rogers, B.A., H.E. Sidjabat, and D.L. Paterson. 2010. *Escherichia coli* O25b-ST131: a pandemic, multiresistant, community-associated strain. *J. Antimicrob. Chemother.* 66:1–14.
5. Banerjee, R., and J.R. Johnson. 2014. A new clone sweeps clean: the enigmatic emergence of *Escherichia coli* ST131. *Antimicrob. Agents Chemother.* 58:4997-5004.
6. Johnson, T.J., J.L. Danzeisen, B. Youmans, K. Case, K. Llop, J. Munoz-Aguayo, C. Flores-Figueroa, M. Aziz, N. Stoesser, and E. Sokurenko. 2016. Separate F-type plasmids have shaped the evolution of the H30 subclone of *Escherichia coli* sequence type 131. *MSphere* 1:e00121–16.
7. Mathers, A.J., G. Peirano, and J.D. Pitout. 2015. *Escherichia coli* ST131: the quintessential example of an international multiresistant high-risk clone. In *Advances in Applied Microbiology*. 90:109–154.
8. Mathers, A.J., G. Peirano, and J.D. Pitout. 2015. The role of epidemic resistance plasmids and international high-risk clones in the spread of multidrug-resistant *Enterobacteriaceae*. *Clin. Microbiol. Rev.* 28:565–591.
9. Totsika, M., S.A. Beatson, S. Sarkar, M.-D. Phan, N.K. Petty, N. Bachmann, M. Szubert, H.E. Sidjabat, D.L. Paterson, and M. Upton. 2011. Insights into a multidrug resistant *Escherichia coli* pathogen of the globally disseminated ST131 lineage: genome analysis and virulence mechanisms. *PloS One* 6:e26578.

10. Woodford, N., A. Carattoli, E. Karisik, A. Underwood, M.J. Ellington, and D.M. Livermore. 2009. Complete nucleotide sequences of plasmids pEK204, pEK499, and pEK516, encoding CTX-M enzymes in three major *Escherichia coli* lineages from the United Kingdom, all belonging to the international O25: H4-ST131 clone. *Antimicrob. Agents Chemother.* 53:4472–4482.
11. Naseer, U., and A. Sundsfjord. 2011. The CTX-M Conundrum: Dissemination of Plasmids and *Escherichia coli* Clones. *Microb. Drug Resist.* 17:83–97.
12. Wright, G.D. 2010. Antibiotic resistance in the environment: a link to the clinic? *Curr. Opin. Microbiol.* 13:589–594.
13. Forsberg, K.J., A. Reyes, B. Wang, E.M. Selleck, M.O.A. Sommer, and G. Dantas. 2012. The Shared Antibiotic Resistome of Soil Bacteria and Human Pathogens. *Science* 337:1107–1111.
14. Wu, J., Y. Huang, D. Rao, Y. Zhang, and K. Yang. 2018. Evidence for environmental dissemination of antibiotic resistance mediated by wild birds. *Front. Microbiol.* 9:745.
15. Petty, N.K., N.L.B. Zakour, M. Stanton-Cook, E. Skippington, M. Totsika, B.M. Forde, M.-D. Phan, D.G. Moriel, K.M. Peters, and M. Davies. 2014. Global dissemination of a multidrug resistant *Escherichia coli* clone. *Proc. Natl. Acad. Sci.* 111:5694–5699.
16. Johnson, J.R., B. Johnston, M.A. Kuskowski, E.V. Sokurenko, and V. Tchesnokova. 2015. Intensity and Mechanisms of Fluoroquinolone Resistance within the H30 and H30Rx Subclones of *Escherichia coli* Sequence Type 131 vs. Other Fluoroquinolone-resistant *E. coli*. *Antimicrob. Agents Chemother.* 59:4471–4480.

17. Johnson, J.R., V. Tchesnokova, B. Johnston, C. Clabots, P.L. Roberts, M. Billig, K. Riddell, P. Rogers, X. Qin, S. Butler-Wu, L.B. Price, M. Aziz, M.-H. Nicolas-Chanoine, C. DebRoy, A. Robicsek, G. Hansen, C. Urban, J. Platell, D.J. Trott, G. Zhanel, S.J. Weissman, B.T. Cookson, F.C. Fang, A.P. Limaye, D. Scholes, S. Chattopadhyay, D.C. Hooper, and E.V. Sokurenko. 2013. Abrupt Emergence of a Single Dominant Multidrug-Resistant Strain of *Escherichia coli*. *J. Infect. Dis.* 207:919–928.
18. Pitout, J.D., and R. DeVinney. 2017. *Escherichia coli* ST131: a multidrug-resistant clone primed for global domination. *F1000Research* 6.
19. Price, L.B., J.R. Johnson, M. Aziz, C. Clabots, B. Johnston, V. Tchesnokova, L. Nordstrom, M. Billig, S. Chattopadhyay, M. Stegger, P.S. Andersen, T. Pearson, K. Riddell, P. Rogers, D. Scholes, B. Kahl, P. Keim, and E.V. Sokurenko. 2013. The Epidemic of Extended-Spectrum- β -Lactamase-Producing *Escherichia coli* ST131 Is Driven by a Single Highly Pathogenic Subclone, H30-Rx. *MBio* 4:e00377-13.
20. Zakour, N.L.B., A.S. Alsheikh-Hussain, M.M. Ashcroft, N.T.K. Nhu, L.W. Roberts, M. Stanton-Cook, M.A. Schembri, and S.A. Beatson. 2016. Sequential acquisition of virulence and fluoroquinolone resistance has shaped the evolution of *Escherichia coli* ST131. *MBio* 7:e00347–16.
21. Coque, T.M., Â. Novais, A. Carattoli, L. Poirel, J. Pitout, L. Peixe, F. Baquero, R. Cantón, and P. Nordmann. 2008. Dissemination of clonally related *Escherichia coli* strains expressing extended-spectrum β -lactamase CTX-M-15. *Emerg. Infect. Dis.* 14:195-200.
22. Nicolas-Chanoine, M.-H., J. Blanco, V. Leflon-Guibout, R. Demarty, M.P. Alonso, M.M. Caniça, Y.-J. Park, J.-P. Lavigne, J. Pitout, and J.R. Johnson. 2008. Intercontinental

- emergence of *Escherichia coli* clone O25: H4-ST131 producing CTX-M-15. *J. Antimicrob. Chemother.* 61:273–281.
23. Peirano, G., A.K. van der Bij, J.L. Freeman, L. Poirel, P. Nordmann, M. Costello, V.L. Tchesnokova, and J.D. Pitout. 2014. Characteristics of *Escherichia coli* sequence type 131 isolates that produce extended-spectrum β -lactamases: global distribution of the H30-Rx sublineage. *Antimicrob. Agents Chemother.* 58:3762–3767.
 24. Matsumura, Y., J.R. Johnson, M. Yamamoto, M. Nagao, M. Tanaka, S. Takakura, S. Ichiyama, K.-S.C.M.S. Group, Y. Matsumura, and M. Yamamoto. 2015. CTX-M-27- and CTX-M-14-producing, ciprofloxacin-resistant *Escherichia coli* of the H30 subclonal group within ST131 drive a Japanese regional ESBL epidemic. *J. Antimicrob. Chemother.* 70:1639–1649.
 25. Ewers, C., M. Grobbel, I. Stamm, P.A. Kopp, I. Diehl, T. Semmler, A. Fruth, J. Beutlich, B. Guerra, and L.H. Wieler. 2010. Emergence of human pandemic O25: H4-ST131 CTX-M-15 extended-spectrum- β -lactamase-producing *Escherichia coli* among companion animals. *J. Antimicrob. Chemother.* 65:651–660.
 26. Jamborova, I., B.D. Johnston, I. Papousek, K. Kachlikova, L. Micenkova, C. Clabots, A. Skalova, K. Chudejova, M. Dolejska, I. Literak, and J.R. Johnson. 2018. Extensive Genetic Commonality among Wildlife, Wastewater, Community, and Nosocomial Isolates of *Escherichia coli* Sequence Type 131 (H30R1 and H30Rx Subclones) That Carry blaCTX-M-27 or blaCTX-M-15. *Antimicrob. Agents Chemother.* 62:e00519-18.
 27. Müller, A., R. Stephan, and M. Nüesch-Inderbinnen. 2016. Distribution of virulence factors in ESBL-producing *Escherichia coli* isolated from the environment, livestock, food and humans. *Sci. Total Environ.* 541:667–672.

28. Overdeest, I., I. Willemsen, M. Rijnsburger, A. Eustace, L. Xu, P. Hawkey, M. Heck, P. Savelkoul, C. Vandenbroucke-Grauls, and K. van der Zwaluw. 2011. Extended-spectrum β -lactamase genes of *Escherichia coli* in chicken meat and humans, The Netherlands. *Emerg. Infect. Dis.* 17:1216-1222.
29. Platell, J.L., J.R. Johnson, R.N. Cobbold, and D.J. Trott. 2011. Multidrug-resistant extraintestinal pathogenic *Escherichia coli* of sequence type ST131 in animals and foods. *Vet. Microbiol.* 153:99–108.
30. Pomba, C., J.D. da Fonseca, B.C. Baptista, J.D. Correia, and L. Martínez-Martínez. 2009. Detection of the pandemic O25-ST131 human virulent *Escherichia coli* CTX-M-15-producing clone harboring the *qnrB2* and *aac (6')-Ib-cr* genes in a dog. *Antimicrob. Agents Chemother.* 53:327–328.
31. Ewers, C., A. Bethe, T. Semmler, S. Guenther, and L.H. Wieler. 2012. Extended-spectrum β -lactamase-producing and AmpC-producing *Escherichia coli* from livestock and companion animals, and their putative impact on public health: a global perspective. *Clin. Microbiol. Infect.* 18:646–655.
32. Bonnedahl, J., and J.D. Järhult. 2014. Antibiotic resistance in wild birds. *Ups. J. Med. Sci.* 119:113–116.
33. Ewers, C., S. Guenther, L.H. Wieler, and P. Schierack. 2009. Mallard ducks – a waterfowl species with high risk of distributing *Escherichia coli* pathogenic for humans. *Environ. Microbiol. Rep.* 1:510–517.
34. Guenther, S., K. Aschenbrenner, I. Stamm, A. Bethe, T. Semmler, A. Stubbe, M. Stubbe, N. Batsajkhan, Y. Glupczynski, and L.H. Wieler. 2012. Comparable high rates of

- extended-spectrum-beta-lactamase-producing *Escherichia coli* in birds of prey from Germany and Mongolia. *PLoS One* 7:e53039.
35. Jamborova, I., M. Dolejska, J. Vojtech, S. Guenther, R. Uricariu, J. Drozdowska, I. Papousek, K. Pasekova, W. Meissner, and J. Hordowski. 2015. Plasmid-mediated resistance to cephalosporins and fluoroquinolones in various *Escherichia coli* sequence types isolated from rooks wintering in Europe. *Appl Env. Microbiol* 81:648–657.
 36. Literak, I., M. Dolejska, D. Janoszowska, J. Hrusakova, W. Meissner, H. Rzycka, S. Bzoma, and A. Cizek. 2010. Antibiotic-resistant *Escherichia coli* bacteria, including strains with genes encoding the extended-spectrum beta-lactamase and QnrS, in waterbirds on the Baltic Sea Coast of Poland. *Appl. Environ. Microbiol.* 76:8126–8134.
 37. Radimersky, T., P. Frolkova, D. Janoszowska, M. Dolejska, P. Svec, E. Roubalova, P. Cikova, A. Cizek, and I. Literak. 2010. Antibiotic resistance in faecal bacteria (*Escherichia coli*, *Enterococcus* spp.) in feral pigeons. *J. Appl. Microbiol.* 109:1687–1695.
 38. Stedt, J., J. Bonnedahl, J. Hernandez, B.J. McMahon, B. Hasan, B. Olsen, M. Drobni, and J. Waldenström. 2014. Antibiotic resistance patterns in *Escherichia coli* from gulls in nine European countries. *Infect. Ecol. Epidemiol.* 4:21565.
 39. Tausova, D., M. Dolejska, A. Cizek, L. Hanusova, J. Hrusakova, O. Svoboda, G. Camlik, and I. Literak. 2012. *Escherichia coli* with extended-spectrum β -lactamase and plasmid-mediated quinolone resistance genes in great cormorants and mallards in Central Europe. *J. Antimicrob. Chemother.* 67:1103–1107.
 40. Vredenburg, J., A.R. Varela, B. Hasan, S. Bertilsson, B. Olsen, C. Narciso-da-Rocha, J. Bonnedahl, J. Stedt, P.M. Da Costa, and C.M. Manaia. 2014. Quinolone-resistant

Escherichia coli isolated from birds of prey in Portugal are genetically distinct from those isolated from water environments and gulls in Portugal, Spain and Sweden. *Environ. Microbiol.* 16:995–1004.

41. Wang, J., Z.-B. Ma, Z.-L. Zeng, X.-W. Yang, Y. Huang, and J.-H. Liu. 2017. The role of wildlife (wild birds) in the global transmission of antimicrobial resistance genes. *Zool. Res.* 38:55-80.
42. Fogarty, L.R., S.K. Haack, M.J. Wolcott, and R.L. Whitman. 2003. Abundance and characteristics of the recreational water quality indicator bacteria *Escherichia coli* and enterococci in gull faeces. *J. Appl. Microbiol.* 94:865–878.
43. Hasan, B., Å. Melhus, L. Sandegren, M. Alam, and B. Olsen. 2014. The gull (*Chroicocephalus brunnicephalus*) as an environmental bioindicator and reservoir for antibiotic resistance on the coastlines of the Bay of Bengal. *Microb. Drug Resist.* 20:466–471.
44. Liubojević, D., V. Radosavljević, and D. Milanov. 2016. The role of gulls (*Laridae*) in the emergence and spreading of antibiotic resistance in the environment. *Worlds Poultr. Sci. J.* 72:853–864.
45. Hernandez, J., J. Bonnedahl, I. Eliasson, A. Wallensten, P. Comstedt, A. Johansson, S. Granholm, Å. Melhus, B. Olsen, and M. Drobni. 2010. Globally disseminated human pathogenic *Escherichia coli* of O25b-ST131 clone, harbouring blaCTX-M-15, found in Glaucous-winged gull at remote Commander Islands, Russia. *Environ. Microbiol. Rep.* 2:329–332.

46. CLSI. 2018. Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals. 4th ed. CLSI supplement VET08. Wayne, PA: Clinical and Laboratory Standards Institute.
47. Dallenne, C., A. Da Costa, D. Decré, C. Favier, and G. Arlet. 2010. Development of a set of multiplex PCR assays for the detection of genes encoding important β -lactamases in Enterobacteriaceae. *J. Antimicrob. Chemother.* 65:490–495.
48. Pérez-Pérez, F.J., and N.D. Hanson. 2002. Detection of plasmid-mediated AmpC β -lactamase genes in clinical isolates by using multiplex PCR. *J. Clin. Microbiol.* 40:2153–2162.
49. Park, C.H., A. Robicsek, G.A. Jacoby, D. Sahm, and D.C. Hooper. 2006. Prevalence in the United States of aac (6′)-Ib-cr encoding a ciprofloxacin-modifying enzyme. *Antimicrob. Agents Chemother.* 50:3953–3955.
50. Robicsek, A., J. Strahilevitz, D.F. Sahm, G.A. Jacoby, and D.C. Hooper. 2006. qnr prevalence in ceftazidime-resistant Enterobacteriaceae isolates from the United States. *Antimicrob. Agents Chemother.* 50:2872–2874.
51. Poirel, L., E. Lagrutta, P. Taylor, J. Pham, and P. Nordmann. 2010. Emergence of metallo- β -lactamase NDM-1-producing multidrug-resistant *Escherichia coli* in Australia. *Antimicrob. Agents Chemother.* 54:4914–4916.
52. Liu, Y.-Y., Y. Wang, T.R. Walsh, L.-X. Yi, R. Zhang, J. Spencer, Y. Doi, G. Tian, B. Dong, and X. Huang. 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect. Dis.* 16:161–168.

53. Rebelo, A.R., V. Bortolaia, J.S. Kjeldgaard, S.K. Pedersen, P. Leekitcharoenphon, I.M. Hansen, B. Guerra, B. Malorny, M. Borowiak, J.A. Hammerl, A. Battisti, A. Franco, P. Alba, A. Perrin-Guyomard, S.A. Granier, C. De Frutos Escobar, S. Malhotra-Kumar, L. Villa, A. Carattoli, and R.S. Hendriksen. 2018. Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Eurosurveillance* 23.6.
54. Paton, A.W., and J.C. Paton. 1998. Detection and Characterization of Shiga Toxigenic *Escherichia coli* by Using Multiplex PCR Assays for *stx 1*, *stx 2*, *eaeA*, Enterohemorrhagic *E. coli hlyA*, *rfb O111*, and *rfb O157*. *J. Clin. Microbiol.* 36:598–602.
55. Vidal, M., E. Kruger, C. Durán, R. Lagos, M. Levine, V. Prado, C. Toro, and R. Vidal. 2005. Single multiplex PCR assay to identify simultaneously the six categories of diarrheagenic *Escherichia coli* associated with enteric infections. *J. Clin. Microbiol.* 43:5362–5365.
56. Blanco, M., J.E. Blanco, E. Rodriguez, I. Abalia, M.P. Alonso, and J. Blanco. 1997. Detection of virulence genes in uropathogenic *Escherichia coli* by polymerase chain reaction (PCR): comparison with results obtained using phenotypic methods. *J. Microbiol. Methods* 31:37–43.
57. Clermont, O., S. Bonacorsi, and E. Bingen. 2000. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl. Environ. Microbiol.* 66:4555–4558.
58. Clermont, O., J.K. Christenson, E. Denamur, and D.M. Gordon. 2013. The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ. Microbiol. Rep.* 5:58–65.

59. Clermont, O., M. Lavollay, S. Vimont, C. Deschamps, C. Forestier, C. Branger, E. Denamur, and G. Arlet. 2008. The CTX-M-15-producing *Escherichia coli* diffusing clone belongs to a highly virulent B2 phylogenetic subgroup. *J. Antimicrob. Chemother.* 61:1024–1028.
60. Clermont, O., H. Dhanji, M. Upton, T. Gibreel, A. Fox, D. Boyd, M.R. Mulvey, P. Nordmann, E. Ruppé, and J.L. Sarthou. 2009. Rapid detection of the O25b-ST131 clone of *Escherichia coli* encompassing the CTX-M-15-producing strains. *J. Antimicrob. Chemother.* 64:274–277.
61. Matsumura, Y., J.D. Pitout, G. Peirano, R. DeVinney, T. Noguchi, M. Yamamoto, R. Gomi, T. Matsuda, S. Nakano, and M. Nagao. 2017. Rapid identification of different *Escherichia coli* ST131 clades. *Antimicrob. Agents Chemother.* 61:e00179-17.
62. Oliver, A., L.M. Weigel, J.K. Rasheed, J.E. McGowan, P. Raney, and F.C. Tenover. 2002. Mechanisms of decreased susceptibility to cefpodoxime in *Escherichia coli*. *Antimicrob. Agents Chemother.* 46:3829–3836.
63. Carattoli, A., A. Bertini, L. Villa, V. Falbo, K.L. Hopkins, and E.J. Threlfall. 2005. Identification of plasmids by PCR-based replicon typing. *J. Microbiol. Methods* 63:219–228.
64. Johnson, J.R., and A.L. Stell. 2000. Extended virulence genotypes of *Escherichia coli* strains from patients with urosepsis in relation to phylogeny and host compromise. *J. Infect. Dis.* 181:261–272.
65. Picard, B., J.S. Garcia, S. Gouriou, P. Duriez, N. Brahimi, E. Bingen, J. Elion, and E. Denamur. 1999. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect. Immun.* 67:546–553.

66. Báez, J., M. Hernández-García, C. Guamparito, S. Díaz, A. Olave, K. Guerrero, R. Cantón, F. Baquero, J. Gahona, N. Valenzuela, R. del Campo, and J. Silva. 2014. Molecular Characterization and Genetic Diversity of ESBL-Producing *Escherichia coli* Colonizing the Migratory Franklin's Gulls (*Leucophaeus pipixcan*) in Antofagasta, North of Chile. *Microb. Drug Resist.* 21:111–116.
67. Bonnedahl, J., M. Drobni, M. Gauthier-Clerc, J. Hernandez, S. Granholm, Y. Kayser, Å. Melhus, G. Kahlmeter, J. Waldenström, and A. Johansson. 2009. Dissemination of *Escherichia coli* with CTX-M type ESBL between humans and yellow-legged gulls in the south of France. *PloS One* 4:e5958.
68. Dolejska, M., A. Cizek, and I. Literak. 2007. High prevalence of antimicrobial-resistant genes and integrons in *Escherichia coli* isolates from black-headed gulls in the Czech Republic. *J. Appl. Microbiol.* 103:11–19.
69. Poeta, P., H. Radhouani, G. Igrejas, A. Gonçalves, C. Carvalho, J. Rodrigues, L. Vinué, S. Somalo, and C. Torres. 2008. Seagulls of the Berlengas natural reserve of Portugal as carriers of fecal *Escherichia coli* harboring CTX-M and TEM extended-spectrum beta-lactamases. *Appl. Environ. Microbiol.* 74:7439–7441.
70. Radhouani, H., P. Poeta, G. Igrejas, A. Goncalves, L. Vinue, and C. Torres. 2009. Antimicrobial resistance and phylogenetic groups in isolates of *Escherichia coli* from seagulls at the Berlengas nature reserve. *Vet. Rec.* 165:138-142.
71. Simões, R.R., L. Poirel, P.M. Da Costa, and P. Nordmann. 2010. Seagulls and beaches as reservoirs for multidrug-resistant *Escherichia coli*. *Emerg. Infect. Dis.* 16:110-112.

72. Stedt, J., J. Bonnedahl, J. Hernandez, J. Waldenström, B.J. McMahon, C. Tolf, B. Olsen, and M. Drobni. 2015. Carriage of CTX-M type extended spectrum β -lactamases (ESBLs) in gulls across Europe. *Acta Vet. Scand.* 57:74.
73. Poirel, L., A. Potron, C. De La Cuesta, T. Cleary, P. Nordmann, and L.S. Munoz-Price. 2012. Wild coastline birds as reservoirs of broad-spectrum B-lactamase-producing Enterobacteriaceae, Miami Beach. *Antimicrob. Agents Chemother.* 56:2756-2758.
74. Veldman, K., P. van Tulden, A. Kant, J. Testerink, and D. Mevius. 2013. Characteristics of cefotaxime resistant *E. coli* from wild birds in The Netherlands. *Appl. Environ. Microbiol.* 79:7556-7561.
75. Vergara, A., C. Pitart, T. Montalvo, I. Roca, S. Sabaté, J.C. Hurtado, R. Planell, F. Marco, B. Ramírez, V. Peracho, M. de Simón, and J. Vila. 2017. Prevalence of Extended-Spectrum- β -Lactamase- and/or Carbapenemase-Producing *Escherichia coli* Isolated from Yellow-Legged Gulls from Barcelona, Spain. *Antimicrob. Agents Chemother.* 61:e02071-16.
76. Vittecoq, M., C. Laurens, L. Brazier, P. Durand, E. Elguero, A. Arnal, F. Thomas, S. Aberkane, N. Renaud, F. Prugnolle, J. Solassol, H. Jean-Pierre, S. Godreuil, and F. Renaud. 2017. VIM-1 carbapenemase-producing *Escherichia coli* in gulls from southern France. *Ecol. Evol.* 7:1224–1232.
77. Mukerji, S., M. Stegger, A.V. Truswell, T. Laird, D. Jordan, R.J. Abraham, A. Harb, M. Barton, M. O’Dea, and S. Abraham. 2019. Resistance to critically important antimicrobials in Australian silver gulls (*Chroicocephalus novaehollandiae*) and evidence of anthropogenic origins. *J. Antimicrob. Chemother.* 74:2566-2574.

78. Allen, H.K., J. Donato, H.H. Wang, K.A. Cloud-Hansen, J. Davies, and J. Handelsman. 2010. Call of the wild: antibiotic resistance genes in natural environments. *Nat. Rev. Microbiol.* 8:251–259.
79. Guenther, S., C. Ewers, and L.H. Wieler. 2011. Extended-spectrum beta-lactamases producing *E. coli* in wildlife, yet another form of environmental pollution? *Front. Microbiol.* 2:246.
80. Skurnik, D., R. Ruimy, A. Andremont, C. Amorin, P. Rouquet, B. Picard, and E. Denamur. 2006. Effect of human vicinity on antimicrobial resistance and integrons in animal faecal *Escherichia coli*. *J. Antimicrob. Chemother.* 57:1215–1219.
81. Alroy, K., and J.C. Ellis. 2011. Pilot study of antimicrobial-resistant *Escherichia coli* in herring gulls (*Larus argentatus*) and wastewater in the northeastern United States. *J. Zoo Wildl. Med. Am. Assoc. Zoo Vet.* 42:160–163.
82. Nelson, M., S.H. Jones, C. Edwards, and J.C. Ellis. 2008. Characterization of *Escherichia coli* populations from gulls, landfill trash, and wastewater using ribotyping. *Dis. Aquat. Organ.* 81:53–63.
83. Ruzauskas, M., and L. Vaskeviciute. 2016. Detection of the *mcr-1* gene in *Escherichia coli* prevalent in the migratory bird species *Larus argentatus*. *J. Antimicrob. Chemother.* 71:2333–2334.
84. Dolejska, M., B. Bierošová, L. Kohoutova, I. Literak, and A. Čížek. 2009. Antibiotic-resistant *Salmonella* and *Escherichia coli* isolates with integrons and extended-spectrum beta-lactamases in surface water and sympatric black-headed gulls. *J. Appl. Microbiol.* 106:1941–1950.

85. Moura, A., S. Araújo, M.S. Alves, I. Henriques, A. Pereira, and A. Correia. 2014. The contribution of *Escherichia coli* from human and animal sources to the integron gene pool in coastal waters. *Front. Microbiol.* 5:419.
86. Bonnedahl, J., P. Drobni, A. Johansson, J. Hernandez, Å. Melhus, J. Stedt, B. Olsen, and M. Drobni. 2010. Characterization, and comparison, of human clinical and black-headed gull (*Larus ridibundus*) extended-spectrum β -lactamase-producing bacterial isolates from Kalmar, on the southeast coast of Sweden. *J. Antimicrob. Chemother.* 65:1939–1944.
87. Hernandez, J., A. Johansson, J. Stedt, S. Bengtsson, A. Porczak, S. Granholm, D. González-Acuña, B. Olsen, J. Bonnedahl, and M. Drobni. 2013. Characterization and comparison of extended-spectrum β -lactamase (ESBL) resistance genotypes and population structure of *Escherichia coli* isolated from Franklin's gulls (*Leucophaeus pipixcan*) and humans in Chile. *PLoS One* 8:e76150.
88. Atterby, C., A.M. Ramey, G.G. Hall, J. Järhult, S. Börjesson, and J. Bonnedahl. 2016. Increased prevalence of antibiotic-resistant *E. coli* in gulls sampled in Southcentral Alaska is associated with urban environments. *Infect. Ecol. Epidemiol.* 6:32334.
89. Matsumura, Y., J.D. Pitout, R. Gomi, T. Matsuda, T. Noguchi, M. Yamamoto, G. Peirano, R. DeVinney, P.A. Bradford, and M.R. Motyl. 2016. Global *Escherichia coli* sequence type 131 clade with *bla*CTX-M-27 gene. *Emerg. Infect. Dis.* 22:1900-1907.
90. Sun, J., X. Liao, L. Li, Y. Yang, B. Liu, Y. Wang, H. Zhu, L. Zhu, C. Tu, and Y. Liu. 2011. Emergence of pandemic B2-O25b-ST131 *Escherichia coli* harbouring *bla*CTX-M-3, *bla*CTX-M-27 and *qnrS1* genes. *J. Med. Microbiol.* 60:1711–1712.

91. Birgy, A., P. Bidet, C. Levy, E. Sobral, R. Cohen, and S. Bonacorsi. 2017. CTX-M-27–Producing *Escherichia coli* of Sequence Type 131 and Clade C1-M27, France. *Emerg. Infect. Dis.* 23:885.
92. Ghosh, H., S. Doijad, L. Falgenhauer, M. Fritzenwanker, C. Imirzalioglu, and T. Chakraborty. 2017. blaCTX-M-27–Encoding *Escherichia coli* Sequence Type 131 Lineage C1-M27 Clone in Clinical Isolates, Germany. *Emerg. Infect. Dis.* 23:1754-1756.
93. Merino, I., M. Hernández-García, M.-C. Turrientes, B. Pérez-Viso, N. López-Fresneña, C. Diaz-Agero, F. Maechler, C. Fankhauser-Rodriguez, A. Kola, and J. Schrenzel. 2018. Emergence of ESBL-producing *Escherichia coli* ST131-C1-M27 clade colonizing patients in Europe. *J. Antimicrob. Chemother.* 73:2973-2980.
94. Rodrigues, C., E. Machado, S. Fernandes, L. Peixe, and Â. Novais. 2016. An update on faecal carriage of ESBL-producing Enterobacteriaceae by Portuguese healthy humans: detection of the H 30 subclone of B2-ST131 *Escherichia coli* producing CTX-M-27. *J. Antimicrob. Chemother.* 71:1120–1122.
95. Bortolami, A., F. Zendri, E.I. Maciuca, A. Wattret, C. Ellis, V. Schmidt, G. Pinchbeck, and D. Timofte. 2019. Diversity, Virulence, and Clinical Significance of Extended-Spectrum β -Lactamase- and pAmpC-Producing *Escherichia coli* From Companion Animals. *Front. Microbiol.* 10:1260.
96. Harada, K., Y. Nakai, and Y. Kataoka. 2012. Mechanisms of resistance to cephalosporin and emergence of O25b-ST131 clone harboring CTX-M-27 β -lactamase in extraintestinal pathogenic *Escherichia coli* from dogs and cats in Japan. *Microbiol. Immunol.* 56:480–485.

97. Kawamura, K., T. Sugawara, N. Matsuo, K. Hayashi, C. Norizuki, K. Tamai, T. Kondo, and Y. Arakawa. 2017. Spread of CTX-Type Extended-Spectrum β -Lactamase-Producing *Escherichia coli* Isolates of Epidemic Clone B2-O25-ST131 Among Dogs and Cats in Japan. *Microb. Drug Resist.* 23:1059–1066.
98. Maeyama, Y., Y. Taniguchi, W. Hayashi, Y. Ohsaki, S. Osaka, S. Koide, K. Tamai, Y. Nagano, Y. Arakawa, and N. Nagano. 2018. Prevalence of ESBL/AmpC genes and specific clones among the third-generation cephalosporin-resistant Enterobacteriaceae from canine and feline clinical specimens in Japan. *Vet. Microbiol.* 216:183–189.
99. Zogg, A.L., K. Zurfluh, S. Schmitt, M. Nüesch-Inderbinnen, and R. Stephan. 2018. Antimicrobial resistance, multilocus sequence types and virulence profiles of ESBL producing and non-ESBL producing uropathogenic *Escherichia coli* isolated from cats and dogs in Switzerland. *Vet. Microbiol.* 216:79–84.
100. Zurfluh, K., M. Nüesch-Inderbinnen, R. Stephan, and H. Hächler. 2013. Higher-generation cephalosporin-resistant *Escherichia coli* in feral birds in Switzerland. *Int. J. Antimicrob. Agents* 41:296–297.
101. Gomi, R., T. Matsuda, Y. Matsumura, M. Yamamoto, M. Tanaka, S. Ichiyama, and M. Yoneda. 2017. Occurrence of clinically important lineages, including the ST131 C1-M27 subclone, among extended-spectrum β -lactamase-producing *Escherichia coli* in wastewater. *Antimicrob. Agents Chemother.* 61:e00564-17.
102. Kittinger, C., M. Lipp, B. Folli, A. Kirschner, R. Baumert, H. Galler, A.J. Grisold, J. Luxner, M. Weissenbacher, and A.H. Farnleitner. 2016. Enterobacteriaceae isolated from the River Danube: Antibiotic resistances, with a focus on the presence of ESBL and carbapenemases. *PloS One* 11:e0165820.

103. Zurfluh, K., H. Hächler, M. Nüesch-Inderbinen, and R. Stephan. 2013. Characteristics of extended-spectrum beta-lactamase (ESBL)- and carbapenemase-producing Enterobacteriaceae isolated from rivers and lakes in Switzerland. *Appl. Environ. Microbiol.* **79**:3021-3026.
104. Amos, G.C.A., P.M. Hawkey, W.H. Gaze, and E.M. Wellington. 2014. Waste water effluent contributes to the dissemination of CTX-M-15 in the natural environment. *J. Antimicrob. Chemother.* **69**:1785–1791.
105. Dhanji, H., N.M. Murphy, C. Akhigbe, M. Doumith, R. Hope, D.M. Livermore, and N. Woodford. 2011. Isolation of fluoroquinolone-resistant O25b:H4-ST131 *Escherichia coli* with CTX-M-14 extended-spectrum β -lactamase from UK river water. *J. Antimicrob. Chemother.* **66**:512–516.
106. Leonard, A.F., L. Zhang, A.J. Balfour, R. Garside, P.M. Hawkey, A.K. Murray, O.C. Ukoumunne, and W.H. Gaze. 2018. Exposure to and colonisation by antibiotic-resistant *E. coli* in UK coastal water users: Environmental surveillance, exposure assessment, and epidemiological study (Beach Bum Survey). *Environ. Int.* **114**:326–333.
107. Platell, J.L., R.N. Cobbold, J.R. Johnson, and D.J. Trott. 2010. Clonal group distribution of fluoroquinolone-resistant *Escherichia coli* among humans and companion animals in Australia. *J. Antimicrob. Chemother.* **65**:1936–1938.